

# Distributed Algorithms 2023

10 Sinkless orientation

# This week's plan

- **Topic:** complexity of *sinkless orientation*
  - task: high-degree nodes must have outdegree  $\geq 1$
  - possible in  $O(\log n)$  rounds, not in  $o(\log n)$  rounds
- **Video:** *why* do we care about this?
  - e.g. hardness of graph coloring
- **Today:** how to *prove* it?
  - round elimination & fixed points

# Sinkless orientation

- **Labels:**  $\{ 0, 1 \}$ 
  - $0$  = *"edge oriented away from the active node"*
  - $1$  = *"edge oriented towards the active node"*
- **Active:**  $[ 0, ?, ? ]$ 
  - *"at least one outgoing edge"*
- **Passive:**  $[ 1, ?, ? ]$ 
  - *"at least one outgoing edge"*

## **Sinkless orientation:** $O, I$

- active:  $[ O, ?, ? ]$
- passive:  $[ I, ?, ? ]$

## **Output problem:** $\{O\}, \{I\}, \{O,I\}$

- active:  $[ \{I\}, ?, ? ]$
- passive:  $[ \{O\}, ?, ? ]$  or  $[ \{O,I\}, ?, ? ]$

## **Maximal problem:** $\{I\}, \{O,I\}$

- active:  $[ \{I\}, \{O,I\}, \{O,I\} ]$
- passive:  $[ \{O,I\}, ?, ? ]$

## **Sinkless orientation:** $O, I$

- active: [  $O, ?, ?$  ]
- passive: [  $I, ?, ?$  ]

## **Output problem:** $\{O\}, \{I\}, \{O,I\}$

- active: [  $\{I\}, ?, ?$  ]
- passive: [  $\{O\}, ?, ?$  ] or [  $\{O,I\}, ?, ?$  ]

## **Maximal problem:** $A, B$

- active: [  $A, B, B$  ]
- passive: [  $B, ?, ?$  ]

# Output problem

- **Labels:** { A, B }
  - A = *"edge oriented away from the active node"*
  - B = *"edge oriented towards the active node"*
- **Active:** [ A, B, B ]
  - *"**exactly** one outgoing edge"*
- **Passive:** [ B, ?, ? ]
  - *"at least one outgoing edge"*

## Starting point: A, B

- active: [ A, B, B ]
- passive: [ B, ?, ? ]

## Output problem: {A}, {B}, {A,B}

- active: [ {B}, ?, ? ]
- passive: ...

## Maximal problem: {B}, {A,B}

- active: [ {B}, {A,B}, {A,B} ]
- passive: [ {A,B}, ?, ? ]

## Starting point: A, B

- active: [ A, B, B ]
- passive: [ B, ?, ? ]

## Output problem: {A}, {B}, {A,B}

- active: [ {B}, ?, ? ]
- passive: ...

## Maximal problem: A, B

- active: [ A, B, B ]
- passive: [ B, ?, ? ]

# Fixed points

- $X = \mathbf{re}(X)$ , and  $X$  is not 0-round solvable
- “ $X$  can be solved 1 round faster than  $X$ ”
  - contradiction
- One of our assumptions fails — which one?

# Fixed points

- $X = \text{re}(X)$ , and  $X$  is not 0-round solvable
- *$X$  cannot be solved in  $o(\log n)$  rounds* in the deterministic PN model
- We can also derive hardness results for deterministic and randomized LOCAL model

# Often used like this

- We are interested in problem  $X$
- Find a suitable *relaxation*  $Y$  of  $X$ 
  - problem  $Y$  is at most as hard as  $X$
  - problem  $Y$  is nontrivial
- Show that  $Y = \text{re}(Y)$  or  $Y = \text{re}(\text{re}(Y))$ 
  - $Y$  cannot be solved fast
  - $X$  cannot be solved fast

# Sinkless and sourceless

- **Labels:**  $\{ 0, 1 \}$ 
  - $0 =$  "edge oriented away from the active node"
  - $1 =$  "edge oriented towards the active node"
- **Active:**  $[ 0, 1, ? ]$ 
  - "at least one outgoing and one incoming edge"
- **Passive:**  $[ 1, 0, ? ]$ 
  - "at least one outgoing and one incoming edge"